

Washington State Department of Health
Disease Summary Table

BOTULISM

Case, Death Rate per 100,000 Population

| Year | Food | Intestinal | Wound | Combined Rate | Deaths | Rate |
|------|------|------------|-------|---------------|--------|------|
| 1985 | 5 | 4 | 0 | 0.2 | 0 | 0.0 |
| 1986 | 2 | 4 | 0 | 0.1 | 0 | 0.0 |
| 1987 | 1 | 1 | 1 | 0.1 | 0 | 0.0 |
| 1988 | 3 | 4 | 0 | 0.2 | 0 | 0.0 |
| 1989 | 10 | 0 | 0 | 0.2 | 0 | 0.0 |
| 1990 | 1 | 0 | 0 | 0.1 | 0 | 0.0 |
| 1991 | 0 | 3 | 0 | 0.1 | 0 | 0.0 |
| 1992 | 0 | 2 | 0 | 0.0 | 0 | 0.0 |
| 1993 | 4 | 5 | 0 | 0.2 | 0 | 0.0 |
| 1994 | 3 | 2 | 0 | 0.1 | 0 | 0.0 |
| 1995 | 4 | 2 | 0 | 0.1 | 0 | 0.0 |
| 1996 | 2 | 0 | 2 | 0.1 | 0 | 0.0 |
| 1997 | 0 | 1 | 2 | 0.1 | 0 | 0.0 |
| 1998 | 2 | 4 | 0 | 0.1 | 0 | 0.0 |
| 1999 | 2 | 4 | 1 | 0.1 | 0 | 0.0 |
| 2000 | 1 | 4 | 0 | 0.1 | 0 | 0.0 |
| 2001 | 1 | 6 | 0 | 0.1 | 0 | 0.0 |
| 2002 | 1 | 1 | 4 | 0.1 | 0 | 0.0 |